The 2011 Montana Youth Risk Behavior Survey

2011 Results and Trends www.opi.mt.gov/yrbs




The Youth Risk Behavior Survey

- Conducted since 1991 by the Centers for Disease Control and Prevention (CDC)
- Measures six topics whose behaviors may contribute to mortality or morbidity.
 - Unintentional injuries and violence
 - Tobacco Use
 - Alcohol and other drug use
 - Sexual behaviors
 - Dietary behaviors
 - Physical inactivity




Percentage of students who ever tried cigarette smoking, even one or two puffs








Percentage of students who ever tried cigarette smoking, even one or two puffs






Percentage of students who smoked a whole cigarette for the first time before age 13 years








Percentage of students who smoked a whole cigarette for the first time before age 13 years






Percentage of students who smoked cigarettes on one or more of the past 30 days








Percentage of students who smoked cigarettes on one or more of the past 30 days





Percentage of students who smoked cigarettes on 20 or more of the past 30 days






*Non-Hispanic.

Missing bars indicate less than 100 students in the subgroup.




Percentage of students who smoked cigarettes on 20 or more of the past 30 days





Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days











Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days






Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days








Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days





Percentage of students who smoked cigarettes on school property on one or more of the past 30 days






Missing bars indicate less than 100 students in the subgroup.




Percentage of students who smoked cigarettes on school property on one or more of the past 30 days





Percentage of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days











Percentage of students who ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days





Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months






Missing bars indicate less than 100 students in the subgroup.




Among students who reported current cigarette use, the percentage who ever tried to quit smoking cigarettes during the past 12 months






Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days








Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days





Percentage of students who used chewing tobacco, snuff, or dip on school property on one or more of the past 30 days











Percentage of students who used chewing tobacco, snuff, or dip on school property on one or more of the past 30 days






Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days









Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days





Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days





*Non-Hispanic.

Missing bars indicate less than 100 students in the subgroup.



Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days

